

$g_{\omega\rho\pi}$ reexamined

M. Lublinsky*

Department of Physics, Technion – Israel Institute of Technology, Haifa 32000, Israel

The transition constant for the hadronic decay $\omega \rightarrow \rho\pi$ is investigated by means of QCD sum rules in external axial field. The obtained value for $g_{\omega\rho\pi}$ is about 16 GeV^{-1} that is in a good agreement with experimental data.

13.25.Jx

I. INTRODUCTION

During last several years, there was a significant increase in the interest in the low energy QCD and in understanding the low energy hadron dynamics. This happened both due to the theoretical progress and to the availability of new experimental data. However, the problem of understanding the low energy properties of hadrons is still far from being solved, due to the need to understand the nonperturbative QCD effects that govern the hadron dynamics.

The QCD sum rules, first suggested by Shifman, Vainshtein and Zakharov [1] (see also Ref. [2] for an extensive review), historically are one of the most effective tools used to study the low energy properties of hadrons. The method was successfully applied to numerous low energy problems, such as the determination of the mass spectrum of the low-lying resonances, of the form-factors etc.

The goal of the present note is to use QCD sum rules in order to calculate the $\omega \rightarrow \rho\pi$ transition constant. Let us recall that this constant is defined [3] by the equation:

$$M_{\omega \rightarrow \rho\pi} = g_{\omega\rho\pi} \epsilon_{ij\alpha\beta} p^i p'^j e_\omega^\alpha e_\rho^\beta \quad (1.1)$$

Here M is the invariant amplitude for the $\omega \rightarrow \rho\pi$ transition, p, e_ω and p', e_ρ denote momenta and polarisations of ω and ρ . Experimental value for $g_{\omega\rho\pi}$ is about $14 \pm 2 \text{ GeV}^{-1}$. Within the framework of QCD sum rules, $g_{\omega\rho\pi}$ was already considered several times [3–8]. The results obtained in Ref. [3–5] varied from 17 GeV^{-1} to 15 GeV^{-1} , that were in an agreement with the experimental data up to 20% - 30% - the usual error of the method. Ref [6] gives for this constant 9 GeV^{-1} that contradicts both the previous calculations and the experimental data. The authors of Ref. [7] present two results. The first one supports calculations of Ref. [6]. In order to obtain the second result they modified sum rules incorporating the ABJ axial anomaly and finally $g_{\omega\rho\pi}$ was estimated as 16 GeV^{-1} . In Ref. [8], the light cone sum rule method was used to determine a wave-function of π meson. The results gave 15 GeV^{-1} for $g_{\omega\rho\pi}$. Some uncertainty in predictions and a significant mismatch between results of Ref. [6] and the other obtained results have drawn us to reconsider $g_{\omega\rho\pi}$.

In this note we calculate $g_{\omega\rho\pi}$ using the QCD sum rules for the polarisation operator in external axial field. This method was first discovered in Ref [9] and since was used

*E-mail: mal@techunix.technion.ac.il

to determine the magnetic moments of baryons [9], axial coupling constants and electromagnetic form-factors of nucleons [10–12] at low momenta transfer, to study $D^0 \rightarrow D\gamma$, Dn transitions [13], etc.

The main advantage of this method is that it permits one to work directly in the region of small momenta transfer (by paying the price of introducing new condensates, see below). Note that this is exactly the case at hand: the masses of ω and ρ are almost equal, hence the momentum of virtual π meson is expected to be small.

The structure of the paper is as follows. In Sec. 2 we shortly remind the reader the basics of QCD sum rules method in external field. Sec. 3 is devoted to the main calculations and in the Sec. 4 we give the conclusions.

II. EXTERNAL FIELD METHOD

Let us review in short the QCD sum rules for polarisation operator in external field.

We start with the polarisation operator:

$$\Pi_A^{\mu\nu}(p, q) = - \int d^4x e^{-ipx} \langle j_\omega^\mu(x) j_\rho^\nu(0) \rangle_A \quad (2.1)$$

Here j_ω^μ, j_ρ^ν are the quark currents that create ω^0 and ρ^0 mesons:

$$\begin{aligned} j_\omega^\mu &= \frac{1}{6}(\bar{u}\gamma^\mu u + \bar{d}\gamma^\mu d) \\ j_\rho^\nu &= \frac{1}{2}(\bar{u}\gamma^\nu u - \bar{d}\gamma^\nu d) \end{aligned} \quad (2.2)$$

The index A means that we consider this correlator in the external axial field $A(x) = Ae^{-iqx}$ that corresponds to the following term in the Lagrangian:

$$L_A = -\frac{1}{2} \int d^4y e^{-iqy} j_5^\lambda A_\lambda \quad (2.3)$$

Here q is the external field momentum and j_5^λ is the quark current that creates pion:

$$j_5^\lambda = \frac{1}{2}(\bar{u}\gamma^\lambda \gamma_5 u - \bar{d}\gamma^\lambda \gamma_5 d) \quad (2.4)$$

The expression (2.1) contains, generally speaking, terms of different powers in A_α . We shall be interested here in the terms linear in A only. They appear both when considering the propagator of the quarks in the external field and when taking into account the interaction of the external field with the quark condensate. Moreover, as it is explained below, it is enough to carry the actual calculations for the case $q_\alpha = 0$, i.e. for the case of constant external field. In the latter case, in the approximation linear in A_α , the quark propagator in the x representation is the following:

$$S_A(x) = -\frac{\hat{x}}{2\pi^2 x^4} - i \frac{(Ax)\hat{x}\gamma_5}{2\pi^2 x^4} \quad (2.5)$$

The terms in the operator expansion which correspond to the nonperturbative interaction of the quarks with the axial field have been obtained in Ref. [10] and have the following form:

$$\begin{aligned} \langle : \bar{\psi}\psi : \rangle_A &= \delta^{ab} \frac{f_\pi^2}{12} \hat{A} \gamma_5 + i \delta^{ab} \frac{\langle : \bar{\psi}\psi : \rangle}{3^2 2^3} (\hat{x} \hat{A} - \hat{A} \hat{x}) \gamma_5 \\ &\quad - \delta^{ab} m_1^2 \frac{f_\pi^2}{6^3} \left(\frac{5}{2} x^3 (A \gamma_5) - (A x) \hat{x} \gamma_5 \right) \end{aligned} \quad (2.6)$$

Here, $f_\pi = 135$ Mev, $\langle : \bar{\psi}\psi : \rangle = -(240 \text{ Mev})^3$, $m_1^2 = 0.2 \text{ GeV}^2$ is the parameter of the matrix element:

$$\langle 0 | g_s \bar{u} \tilde{G}_{\mu\nu}^i (\lambda^i / 2) \gamma_\nu d | \pi \rangle = i m_1^2 f_\pi q_\mu$$

which is defined in Ref. [14], and $\tilde{G}_{\mu\nu}^i$ is the dual tensor of the gluon field strength. By means of expressions (2.5), (2.6), and using the technique developed in Ref. [9,10], we are able to determine the polarisation operator $\Pi_A^{\mu\nu}(p, q)$ in the region $p^2, q^2 \sim -1 \text{ GeV}^2$. The actual calculation is carried through using the Wilson operator product expansion (OPE) and taking into account several first low dimensional operators:

$$\gamma_\mu(x) \gamma_\nu(0) = \sum C_n(x^2) O_n(0),$$

where $C_n(x^2)$ are coefficient functions.

In order to extract pion pole from the amplitude (1.1) we have to manage the case of small, but nonzero q . It was suggested in Ref. [13] that in the case of small but nonzero q , one may perform the calculation of the polarisation operator for $q = 0$ and then, in the final expression, make the substitution:

$$A_\alpha \rightarrow (g_\alpha^\lambda - \frac{q_\alpha q^\lambda}{q^2}) A_\lambda \quad (2.7)$$

The calculation of the polarisation operator (2.1) by means of OPE gives the right-hand, theoretical part of the sum rule. On the other side, the polarisation operator satisfies the dispersion relations, which we shall saturate by the contribution of the intermediate states. The dispersion relations are:

$$\begin{aligned} f(Q^2) &= \int_0^\infty \frac{\rho(s) ds}{(s + Q^2)^2} + \int_0^\infty \frac{\rho_1(s) ds}{s + Q^2} \\ Q^2 &= -p^2 \end{aligned} \quad (2.8)$$

where $\rho(s)$ is the contribution in (2.1) of the diagonal transitions corresponding to the diagram (Fig. 1), $\rho_1(s)$ is the contribution of the non-diagonal transitions (Fig. 2). $f(Q^2)$ denotes the structure function of the polarisation operator $\Pi_A^{\mu\nu}$ that we use for the calculation of $g_{\omega\rho\pi}$.

After saturating (2.8) by resonances, taking into account ω, ρ resonance contributions explicitly, and taking care of higher excited states using continuum model [15], we obtain the phenomenological, left-hand part of the sum rule.

III. SUM RULE

Apply now the ideas discussed above to the $\omega \rightarrow \rho\pi$ transition.

To begin with, consider the phenomenological part. We saturate the currents (2.2), (2.4) by the physical resonances and define the residues:

$$\begin{aligned}\langle 0|j_\omega^\mu|\omega\rangle &= \frac{m_\omega^2}{g_\omega}e_\omega^\mu \\ \langle 0|j_\rho^\nu|\rho\rangle &= \frac{m_\rho^2}{g_\rho}e_\rho^\nu \\ \langle 0|j_5^\lambda|\pi\rangle &= -i\frac{f_\pi}{\sqrt{2}}q^\lambda\end{aligned}\tag{3.1}$$

Here, m_ω, g_ω and m_ρ, g_ρ denote masses and form-factors of ω and ρ mesons. f_π and q are the form-factor of π meson in chiral limit and its momentum.

The linear (in A) part of the polarisation operator $\Pi_A^{\mu\nu}(p, q)$ is proportional to the invariant amplitude (1.1) and is given by:

$$\Pi_A^{\mu\nu}(p, q) = M \frac{\langle 0|j_5^\lambda|\pi\rangle A_\lambda}{q^2} \frac{\langle 0|j_\omega^\mu|\omega\rangle}{(p^2 - m_\omega^2)} \frac{\langle 0|j_\rho^\nu|\rho\rangle}{(p'^2 - m_\rho^2)} + \dots\tag{3.2}$$

$$p' = p + q$$

The ... denote contributions of other excited states and “parasitic” non-diagonal pieces. Taking limit $q \rightarrow 0$ and substituting (1.1) and (3.1), into (3.2) we arrive to the expression for $\Pi^{\mu\nu}(p, q)$:

$$\Pi_A^{\mu\nu}(p, q) = -ig_{\omega\rho\pi} \frac{m_\rho^2 m_\omega^2}{\sqrt{2}g_\rho g_\omega} \frac{f_\pi}{(p^2 - m_\rho^2)^2} \frac{q^\lambda A_\lambda}{q^2} \epsilon_{\mu\nu\alpha\beta} q^\alpha p^\beta\tag{3.3}$$

In order to obtain (3.3) we used the well-known expression

$$\langle e^\alpha e^\beta \rangle = g^{\alpha\beta} - \frac{p^\alpha p^\beta}{p^2}$$

for the density matrix, we took into account $m_\omega \approx m_\rho$, and we have written explicitly the contribution of the $\omega \rightarrow \rho$ transition only. We need to consider the sum rule for the following structure:

$$\frac{q^\lambda A_\lambda}{q^2} \epsilon_{\mu\nu\alpha\beta} q^\alpha p^\beta\tag{3.4}$$

In the subsequent calculations of OPE we keep only the terms that contain the structure (3.4).

Let us now turn to the left, theoretical part. We restrict ourselves to the operators of dimensions $d \leq 5$ only.

Consider the unit operator, first. Its contribution is given by one diagram of zero order in α_s (Fig. 3) and six diagrams of the first order in α_s (Fig. 4).

The explicit calculation of the expression corresponding to the (Fig. 3) gives zero.

Since the contribution of zero order to the unit operator I vanishes, the α_s corrections to this operator are taken into account. We shall do it because the contribution of this operator is potentially the leading one and even α_s corrections due to the unit operator may give large contribution to the sum rule. The α_s corrections are given by diagrams of Fig. 4 and contain one gluon exchange. We calculate the expressions corresponding to all six diagrams (Fig. 4) at zero momenta transfer. Subsequently, we perform the substitution (2.7) and retain only the terms containing the structure (3.4). The contribution of each diagram has the same structure:

$$- \phi_i \frac{g^2}{(4\pi)^4} \log\left[\frac{p^2}{4\pi\mu^2}\right] \frac{q^\lambda A_\lambda}{q^2} \epsilon_{\mu\nu\alpha\beta} q^\alpha p^\beta \quad (3.5)$$

$$\alpha_s = \frac{g^2}{(4\pi)}$$

The obtained numerical factors ϕ_i 's are listed below:

$$\begin{aligned} \phi_a + \phi_b &= -i\frac{16}{3}; & \phi_c &= i\frac{16}{9}; \\ \phi_d &= i\frac{4}{9}; & \phi_e + \phi_f &= i\frac{56}{9}; \end{aligned}$$

The appearance of the log term, of course, does not surprise. What is less obvious is a complete cancellation of all infrared divergencies that occur in the mass-less limit we are working in. This remarkable fact signifies the existence of OPE and the fact that long and short distances are separated indeed. Taking the sum, we obtain that the contribution of the unit operator to OPE is given by:

$$- i \frac{28}{9(4\pi)^3} \alpha_s(p^2) \log\left[\frac{p^2}{4\pi\mu^2}\right] \frac{q^\lambda A_\lambda}{q^2} \epsilon_{\mu\nu\alpha\beta} q^\alpha p^\beta \quad (3.6)$$

Here the running coupling constant $\alpha_s(p^2)$ is given by:

$$\alpha_s(p^2) = \frac{4\pi}{9 \log\left[\frac{p^2}{\Lambda^2}\right]} \quad (3.7)$$

Λ denotes the cutoff of QCD and is taken to be 0.2 GeV.

Next, let us calculate the power corrections to the sum rule. We take into account $\langle : \bar{\psi}\psi : \rangle$ and $\langle : \bar{\psi}\psi : \rangle_A$ condensates only. The contributions due to $\langle : GG : \rangle$, $\langle : GG : \rangle_A$, $\langle : \bar{\psi}G\psi : \rangle$, $\langle : \bar{\psi}G\psi : \rangle_A$ are negligible and the operators with $d \geq 6$ are omitted as well. We work in the leading order in α_s .

The power corrections are given by the diagrams of the Fig. 5, Fig. 6.

Odd number of γ matrices entering into the expression for the diagram Fig. 5 forces it to be zero.

The expression corresponding to the diagram Fig. 6 looks like:

$$\begin{aligned} \langle : \psi^b(0) \bar{\psi}^a(x) : \rangle_A \gamma^\mu S_0(x) \gamma^\nu \delta^{ab} \\ - S_0(x) \gamma^\mu \langle : \psi^a(x) \bar{\psi}^b(0) : \rangle_A \gamma^\nu \delta^{ab} \end{aligned}$$

Here S_0 denote the standard quark propagator in x representation. We use (2.6) for the condensates. Performing the substitution (2.7) and retaining only terms containing the structure (3.4) we obtain the condensate contribution to OPE. It has the form:

$$i \left[\frac{f_\pi^2}{3p^2} - \frac{5}{27} \frac{m_1^2 f_\pi^2}{p^4} \right] \frac{q^\lambda A_\lambda}{q^2} \epsilon_{\mu\nu\alpha\beta} q^\alpha p^\beta \quad (3.8)$$

Now in order to obtain the desired expression for $g_{\omega\rho\pi}$ it is a good time to consider the sum rule. On the left hand side we have the expression (3.3), while (3.6) + (3.8) stands on the right hand side.

The obtained sum rule reads:

$$\begin{aligned} g_{\omega\rho\pi} \frac{m_\rho^2 m_\omega^2}{\sqrt{2} g_\rho g_\omega} \frac{f_\pi}{(p^2 - m_\rho^2)^2} + \dots \\ = \frac{5}{27} \frac{m_1^2 f_\pi^2}{p^4} - \frac{f_\pi^2}{3p^2} + \frac{28}{9(4\pi)^3} \alpha_s(p^2) \log[p^2/4\pi\mu^2] \end{aligned} \quad (3.9)$$

In order to suppress the contributions of higher resonances it is essential to perform the Borel transformation [1]. We introduce the continuum threshold S_0 in a standard way and transfer the continuum contribution into the theoretical part of the sum rule. The integration in the dispersion representation of the unit operator is cut to the integral from 0 to S_0 . (Other diagrams have no imaginary part).

Finally, we take into account the anomalous dimensions of the operators [16] and obtain the following sum rule:

$$\begin{aligned} [g_{\omega\rho\pi} + CM^2] \frac{f_\pi m_\rho^2 m_\omega^2}{\sqrt{2} g_\rho g_\omega} e^{-\frac{m_\rho^2}{M^2}} = \frac{5}{27} m_1^2 f_\pi^2 L^\gamma + \frac{f_\pi^2}{3} M^2 \\ + \frac{28}{9(4\pi)^3} \alpha_s(M^2) M^4 (1 - e^{-S_0/M^2}) \end{aligned} \quad (3.10)$$

The piece proportional to C takes into account the non-diagonal transitions given by Fig. 2. Here

$$L = \frac{\log[M^2/\Lambda^2]}{\log[\mu^2/\Lambda^2]}; \quad \gamma = -4/9; \quad \mu = 0.5 \text{ GeV}.$$

For m_ρ^2 and the form-factors the following values are taken:

$$m_\rho^2 = 0.59 \text{ GeV}^2; \quad \frac{4\pi}{g_\rho} = 0.41; \quad \frac{4\pi}{g_\omega} = 0.046.$$

We analyse the obtained sum rule. We find the region of stability, where the contribution of the continuum is less than 30%. In order to have a right hierarchy of the power corrections we look for a region, where the first term of the right-hand side of (3.10) is also less than 30%.

In order to extract $g_{\omega\rho\pi}$ from the obtained sum rule, we transform the equation (3.10) and plot the function $R(M^2, S_0) = g_{\omega\rho\pi} + CM^2$ for various values of S_0 . The desired value

for $g_{\omega\rho\pi}$ is given by the point where the straight lines asymptotically cross the y -axis (Fig. 7). Investigating the graph Fig. 7 we see that the region of stability begins at $M^2 \simeq 1.2 \text{ GeV}^2$. The determined value for $g_{\omega\rho\pi}$:

$$g_{\omega\rho\pi} \sim 16 \text{ GeV}^{-1}.$$

The inaccuracy of the obtained value is about 10%. The result does not change significantly with the variation of S_0 that implies an insensitivity to the continuum threshold. Note also that the α_s corrections seem to be unimportant and the main contribution to the sum rule comes from the condensate terms.

IV. CONCLUSIONS

In the present work we investigated the constant of the $\omega \rightarrow \rho\pi$ transition by means of the QCD sum rules in external field. As was expected, at small momentum transfer the method of external field works perfectly and provides us with a reliable answer. No instability of the sum rules has occurred. The stability region and the optimal value for the parameter of continuum almost coincide with the values, which were used to determine the mass of ρ meson [1].

The obtained value for $g_{\omega\rho\pi}$ is about 16 GeV^{-1} with the relative error of about 10%. Our result is in a good agreement with the experimental data, confirms the calculations of [3–5,7] and disproves the results of Ref. [6].

ACKNOWLEDGMENTS

I am deeply grateful to my supervisor Prof. Boris Blok for the suggestion of the problem and the permanent guidance during the work.

-
- [1] M.A. Shifman, A.I. Vainshtein, V.I. Zakharov, Nucl. Phys. B **147**, 385, 448 (1979).
 - [2] L.I. Reinders, H.R. Rubinstein, S. Yazaki, Phys. Rep. **127**, (1985).
 - [3] V.L. Eletsky, B.L. Ioffe, Y.I. Kogan, Phys. Lett. B **122**, 423, (1983); V.L. Eletsky, Y.I. Kogan, Yad. Fiz. **39**, 138, (1984).
 - [4] L.I. Reinders, H.R. Rubinstein, S. Yazaki, Nucl. Phys. B **213**, 109, (1983).
 - [5] S.Narison, N.Paver, Z. Phys. C **22**, 69, (1984).
 - [6] V.M. Khathimovsky, Yad. Fiz. **41**, 519, (1985).
 - [7] M.V. Margvelashvili, M.E. Shaposhnikov, Z. Phys. C **38**, 467, (1988).
 - [8] V.M. Braun, I.E. Filyanov, Z. Phys. C **44**, 157, (1989).
 - [9] B.L.Ioffe, A.V.Smilga, JETP Lett. **37**, 298, (1983); Nucl. Phys. B **232**, 109, (1984).
 - [10] V.M. Belyaev, Ya.I. Kogan, JETP Lett. **37**, 611, (1983); Phys. Lett. B **135**, 273, (1984); V.M. Belyaev, B.Y. Blok, Ya.I. Kogan, Yad. Fiz. **41**, 439, (1985).
 - [11] V.A. Nesterenko, A.V. Radyushkin, JETP Lett. B **39**, 576, (1984).
 - [12] V.M. Belyaev, Ya.I. Kogan, IJMP A **8**, 153, (1993).

- [13] V.L. Eletsky, Ya.I. Kogan, Z. Phys. C **28**, 155, (1985)
- [14] V.N. Novikov, M.A. Shifman, A.I. Vainshtein and V.I. Zakharov, Nucl. Phys. B **237**, 525, (1984)
- [15] B.L.Ioffe, Nucl. Phys. B **188**, 317, (1981)
- [16] V.M. Belyaev, B.L.Ioffe, JETP **83**, 876, (1982)

FIG. 1. The $\omega \rightarrow \rho$ transition in the presence of external field.

FIG. 2. The parasitic contributions to the $\omega \rightarrow \rho$ transition.

FIG. 3. Contribution of the unit operator to the polarisation operator, of zero order in α_s .

FIG. 4. Contribution of the unit operator to the polarisation operator, of the first order in α_s .

FIG. 5. $\langle : \bar{\psi} \psi : \rangle$ contribution to the polarisation operator.

FIG. 6. $\langle : \bar{\psi} \psi : \rangle_A$ contribution to the polarisation operator.

FIG. 7. The value for $g_{\omega\rho\pi}$ from the sum rule. The cluster of the six curves corresponds to the values of S_0 from 0 to 2.5. The outstanding curve corresponds to $S_0 \rightarrow \infty$.

This figure "fig1-1.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9608331v1>

This figure "fig2-1.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9608331v1>

This figure "fig1-2.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9608331v1>

This figure "fig2-2.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9608331v1>

This figure "fig1-3.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9608331v1>

This figure "fig2-3.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9608331v1>